

144th Estuaries Open Seminar 第144回 汽水域懇談会

East Asian winter monsoon intensity in the Pleistocene based on seasonal land snail shell stable isotope ratios

カタツムリ殻の季節毎の安定同位体比に基づいた
更新世における東アジア冬季モンスーン強度

日時: 2019年 6月19日(水) 17:00-18:00, 19th June (Wed), 2019

場所: エスチュアリー研究センター
2階セミナー室, Seminar Room 2F, EsReC

Dr. David L. Dettman

University of Arizona, USA

デイビッド デットマン (エスチュアリー研究センター客員研究員)



Monsoon climates are characterized by strong seasonal oscillations between a very wet season and very dry conditions. The study of ancient variation in monsoon climate strength and its impact on regional environments is difficult because geological materials rarely record individual seasons. One solution is to use high-resolution sampling of land snail shells as monitors of wet-season and dry-season differences. We subsampled shells along the spiral growth axis, producing 15 to 35 samples per shell to document maximum and minimum oxygen and carbon stable isotope ratios. The most robust signal in land snail shell oxygen isotope records is connected to evaporation. The arid dry-season leads to significant evaporation in the land snail body water, which elevates the $d_{18}O$ values recorded in the shell. Wet-season oxygen isotope ratios are more negative and can be tied roughly to the $d_{18}O$ of wet season precipitation and summer temperatures. Carbon isotope ratios are tied to the diet of the animals, and reveal the proportion of C_3 and C_4 plants in the landscape.

Larger land snail (> 1.5 cm diameter) taxa have behaviors that can lead to significant evaporation of body water, as they are willing to range away from local water sources. Modern samples show that the total amplitude in oxygen isotope variation in a shell can be related to the intensity and duration of the dry season. This suggests that amplitudes in ancient shells are indicators of the winter monsoon strength. The most negative $d_{18}O$ values are not correlated with the intensity of the summer monsoon, suggesting that $d_{18}O$ values are not good indicators of summer monsoon strength.

Data from one genus of land snail (*Cathaica* species) living on the Chinese loess plateau throughout the late Pleistocene allow us to make some generalizations about monsoon climate in this region. In general, the oxygen isotope amplitude in shells from glacial periods are larger than those from the interglacial periods. This suggests that the East Asian Winter Monsoon was strongest during glacial intervals on the Loess Plateau. The largest $d_{18}O$ amplitude observed was 19.2 ‰. Marine Isotope Stages 2, 6, and 8 have the largest amplitudes and, by inference, the strongest winter monsoons in our record. Older glacial intervals seem to have weaker winter monsoons. However, glacial amplitudes were not always larger, just as climate undergoes significant short-term variation in glacial intervals (such as D-O cycles and Heinrich events) oxygen isotope amplitudes show large amounts of variability in glacial times. In contrast, interglacial intervals tend to have smaller $d_{18}O$ amplitudes, pointing to warmer and wetter winters in interglacial times. But like the glacial data, some interglacial amplitudes are larger than some glacial records. These data show that the differences in high frequency climate variability is as large as the differences between glacial and interglacial climates.

お問い合わせ: 島根大学 研究・学術情報機構 エスチュアリー研究センター
センター長・教授 齋藤 文紀 TEL 0852-32-6037